

AD-A058 941

FLORIDA STATE UNIV TALLAHASSEE DEPT OF STATISTICS F/G 12/1  
ON THE NEGATIVE BINOMIAL CONVERGENCE IN A CLASS OF M-DIMENSIONAL--ETC(U)  
JUL 78 H LACAYO, N A LANGBERG AFOSR-74-2581

UNCLASSIFIED

FSU-STATISTICS-M464

AFOSR-TR-78-1232

NL

1 OF 1  
ADA  
058941



END  
DATE  
FILMED  
11-78  
DOC

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

LEVEL 2

## REPORT DOCUMENTATION PAGE

READ INSTRUCTIONS  
BEFORE COMPLETING FORM

1. REPORT NUMBER AFOSR-TR-78-1232		2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle)  ON THE NEGATIVE BINOMIAL CONVERGENCE IN A CLASS OF M-DIMENSIONAL SIMPLE EPIDEMICS		5. TYPE OF REPORT & PERIOD COVERED Interim	
7. AUTHOR(s)  H. Lacayo and Naftali A. Langberg		6. PERFORMING ORG. REPORT NUMBER FSU No. M464	
9. PERFORMING ORGANIZATION NAME AND ADDRESS The Florida State University Department of Statistics Tallahassee, Florida 32306		8. CONTRACT OR GRANT NUMBER(s) AFOSR 76-3109 ✓	
11. CONTROLLING OFFICE NAME AND ADDRESS Air Force Office of Scientific Research/NM Bolling AFB, Washington, DC 20332		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 61102F 2304/A5	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE July 1978	
		13. NUMBER OF PAGES 10	
		15. SECURITY CLASS. (of this report) UNCLASSIFIED	
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release; distribution unlimited.			
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)			
18. SUPPLEMENTARY NOTES			
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)  Exponential distributions, convergence in law, stochastic processes, m-dimensional simple epidemics, negative binomial distributions.			
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)  We consider a population which is exposed to m infections, and consist initially of N susceptibles. At each point in time at most one susceptible becomes infective, and only from one cause. This m-dimensional simple epidemic is a stochastic process, $(X_{sub N,1}(t), \dots, S_{N,m-sub N,1,m}(t))$ , with components counting the number of infectives from the respective causes at time t. We show that if the transition rates of cause 1 through m at time t are given by (continued on reverse side)			

DDC  
SEP 20 1978  
F

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

AD A058941

DDC FILE COPY

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

## 20. Abstract

$\alpha_i X_{N,i}(t) [1 - \frac{1}{N} \sum_{i=1}^m (X_{N,i}(t) - X_{N,i}(0))]$ ,  $i = 1, \dots, m$ ,  $\alpha_1, \dots, \alpha_m > 0$  and  
if  $\lim_{N \rightarrow \infty} X_{N,i}(0) = b_i \in \{1, 2, \dots\}$ , then  $(X_{N,1}(t), \dots, X_{N,m}(t))$  converges as  
 $N \rightarrow \infty$  to a random vector with independent negative binomial components.

ACCESSION for	
NTIS	White Section <input checked="" type="checkbox"/>
DDC	B-11 5a 01 <input type="checkbox"/>
UNANIMOUS	<input type="checkbox"/>
BY	
DISTRIBUTION/AVAILABILITY CODES	
ALL	and/or SPECIAL
A	

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)



⑥ ON THE NEGATIVE BINOMIAL CONVERGENCE  
IN A CLASS OF M-DIMENSIONAL SIMPLE EPIDEMICS,

by ⑨ Interim technical rept.

⑩ H. /Lacayo and Naftali A./Langberg

FSU Statistics Report M464  
AFOSR Technical Report No. 84  
AFOSRN Technical Report No. 9

⑪ Jul 78

⑫ 16p.

July, 1978  
The Florida State University  
Department of Statistics  
Tallahassee, Florida 32306

⑭ FSU-STATISTICS-M464, TR-84-AFOSR

Research sponsored by the Air Force Office of Scientific Research, AFSC,  
USAF, under Grants AFOSR 74-2581D and AFOSR 76-3109.

⑮ ✓ AFOSR-74-2581,  
✓ AFOSR-76-3109

⑯ 2304

⑰ A5

⑱ AFOSR

⑲ TR-78-1232

400 277

78 09 05 067

Approved for public release;  
distribution unlimited.

JLB

ON THE NEGATIVE IONOSPHERIC CONDUCTANCE  
IN A CLASS OF H-IONOSPHERIC STORMS

by

H. JACOBY and H. J. LAMPERT

AFOSR Technical Report No. 84  
AFOSR Technical Report No. 84  
AFOSR Technical Report No. 84

July, 1975  
The Florida State University  
Department of Statistics  
Tallahassee, Florida 32306

Research sponsored by the Air Force Office of Scientific Research, AFOSR,  
under Grant AFOSR 74-2581D and AFOSR 75-2100

**AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFOSR)**  
**NOTICE OF TRANSMITTAL TO DDC**  
This technical report has been reviewed and is  
approved for public release IAW AFR 190-12 (7b).  
Distribution is unlimited.  
**A. D. BLOSE**  
Technical Information Officer

Approved for public release  
Distribution unlimited

On the Negative Binomial Convergence  
in a Class of m-Dimensional Simple Epidemics

by

H. Lacayo and Naftali A. Langberg

ABSTRACT

We consider a population which is exposed to  $m$  infections, and consist initially of  $N$  susceptibles. At each point in time at most one susceptible becomes infective, and only from one cause. This  $m$ -dimensional simple epidemic is a stochastic process,  $(X_{N,1}(t), \dots, X_{N,m}(t))$ , with components counting the number of infectives from the respective causes at time  $t$ . We show that if the transition rates of cause 1 through  $m$  at time  $t$  are given by  $\alpha_i X_{N,i}(t) [1 - \frac{1}{N} \sum_{i=1}^m (X_{N,i}(t) - X_{N,i}(0))]$   $i = 1, \dots, m$ ,  $\alpha_1, \dots, \alpha_m > 0$  and if  $\lim_{N \rightarrow \infty} X_{N,i}(0) = b_i \in \{1, 2, \dots\}$ , then  $(X_{N,1}(t), \dots, X_{N,m}(t))$  converges as  $N \rightarrow \infty$  to a random vector with independent negative binomial components.

**Key Words:** Exponential distributions, convergence in law, stochastic processes,  $m$ -dimensional simple epidemics, negative binomial distributions.

78 09 05 067



## 1. Introduction and Summary

Introduction.

We consider a population which is exposed to  $m$  infections and consists initially of  $N$  susceptibles and  $b_{N,i}$  infectives from cause  $i$ ,  $i = 1, \dots, m$ . At each point in time at most one susceptible becomes an infective, and only from one cause. Once an individual enters the infective state he remains there and cannot be a carrier of any other infection. This  $m$ -dimensional simple epidemic is a stochastic process,  $(X_{N,1}(t), \dots, X_{N,m}(t))$ , with components counting the number of infectives from the respective causes at time  $t$ . We show that if the transition rates of cause 1 through  $m$  at time  $t$  are given by:

$$\alpha_i X_{N,i}(t) [1 - \frac{1}{N} \sum_{i=1}^m (X_{N,i}(t) - b_{N,i})], \quad i = 1, \dots, m, \quad \alpha_1, \dots, \alpha_m > 0, \quad (1.1)$$

and if

$$\lim_{N \rightarrow \infty} b_{N,i} = b_i \in \{1, 2, \dots\}, \quad i = 1, \dots, m \quad (1.2)$$

then for every positive real numbers  $t$  and  $\beta$ , and  $m$  nonnegative integers  $k_1, \dots, k_m$ ,

$$\lim_{N \rightarrow \infty} P\{X_{N,i}(t) \geq k_i + b_i, \quad i = 1, \dots, m\} = \prod_{i=1}^m P\{X_i(t) \geq k_i + b_i\} \quad (1.3)$$

and

$$\lim_{N \rightarrow \infty} E(X_{N,i}(t))^\beta = E(X_i(t))^\beta, \quad i = 1, \dots, m, \quad (1.4)$$

where  $X_1(t), \dots, X_m(t)$  are negative binomial random variables with

respective parameters  $b_i$  and  $e^{-\alpha_i t}$ ,  $i = 1, \dots, m$ . The only previous work in the area of  $m$ -dimensional simple epidemics is the one by Billard, Lacayo and Langberg (1978), who proved (1.3) and (1.4) with the additional assumption that  $\alpha_1 = \alpha_2 = \dots = \alpha_m$ . In that case the interinfection random times are independent of the infection causes. Generally, as will be pointed out in Section 2 this property does not hold. The proof of (1.3) in the cited reference depends on the special structure of the interinfection times. To obtain (1.3) and (1.4) we have approached the problem from an alternate viewpoint.

#### Summary.

In Section 2 we present a rigorous definition of an  $m$ -dimensional simple epidemic, and describe the ones that are the subject of our analysis. Statements (1.3) and (1.4) are proven in Section 3. In this section we present an  $m$ -dimensional simple epidemic in a population consisting initially of infinitely many susceptibles. This theoretical simple epidemic is instrumental in the proofs of (1.3) and (1.4).



## 2. The m-dimensional Simple Epidemic

An m-dimensional simple epidemic in a population consisting initially of N susceptibles and  $b_{N,i}$  infectives from cause i,  $i = 1, \dots, m$  can be described by N bivariate random vectors  $(T_{N,1}, \xi_{N,1}), \dots, (T_{N,N}, \xi_{N,N})$ .  $T_{N,1}, \dots, T_{N,N}$  are the random interinfection times, and  $\xi_{N,1}, \dots, \xi_{N,N}$  are discrete random variables with values in  $\{1, \dots, m\}$  designating the respective infection causes. Let  $S_{N,k}$  be equal to  $\sum_{j=1}^k T_{N,j}$ ,  $k = 1, \dots, N$ . The finite dimensional joint distributions of  $(X_{N,1}(t), \dots, X_{N,m}(t))$  are determined by  $(T_{N,1}, \xi_{N,1}), \dots, (T_{N,N}, \xi_{N,N})$ , through the following set equalities:

$$X_{N,i}(t) = b_{N,i} + k = \bigcup_{r=k}^N (S_{N,r} \leq t < S_{N,r+1}, \sum_{q=1}^r I(\xi_{N,q} = i) = k), t > 0 \quad (2.1)$$

$k = 0, \dots, N$ , and  $i = 1, \dots, m$ . ( $S_{N,0} = 0$ ,  $S_{N,N+1} = \infty$ ).

We assume throughout that  $(T_{N,1}, \xi_{N,1}), \dots, (T_{N,N}, \xi_{N,N})$  are determined by (2.2) and (2.3).

$$P\{\xi_{N,j} = i_j, j = 1, \dots, N\} = \left[ \prod_{i=1}^m \alpha_i \prod_{j=0}^{r_i-1} (b_{N,i} + j) \right] \left[ \prod_{k=1}^N \left( \sum_{i=1}^m \alpha_i (b_{N,i} + \sum_{q=1}^{k-1} I(\xi_{N,q} = i)) \right)^{-1} \right], \quad (2.2)$$

where  $r_i = \sum_{q=1}^N I(\xi_{N,q} = i)$ ,  $i = 1, \dots, m$ .

$$P\{T_{N,k} | \xi_{N,1}, \dots, \xi_{N,k-1} > t_k, k = 1, \dots, N\} = e^{-\sum_{j=1}^N \mu_j t_j} \quad (2.3)$$

where  $t_1, \dots, t_n$  are positive real numbers, and  $\mu_j$  is equal to  $\frac{N-j+1}{N} \sum_{i=1}^m \alpha_i (b_i + \sum_{q=1}^{j-1} I(\xi_q = i))$ , for  $j = 1, \dots, N$ .

In the particular case discussed in Billard, Lacayo and Langberg (1978), condition (2.3) reduces to independent exponential interinfection times

with rates equal to  $\alpha \frac{N - k + 1}{N} (\sum_{i=1}^m b_{N,i} + k - 1)$ ,  $k = 1, \dots, N$ , independent of  $\xi_{N,1}, \dots, \xi_{N,N}$ . In addition equation (2.2) simplifies to  $[\prod_{i=1}^m \prod_{j=0}^{i-1} (b_{N,i} + j)] [\prod_{k=1}^N (\sum_{i=1}^m b_{N,i} + k - 1)]^{-1}$ . Finally we note that if (2.2) and (2.3) hold, then the  $m$ -dimensional simple epidemic satisfies (1.1). Conversely, if (1.1) is satisfied, then equations (2.2) and (2.3) follow by the inherited Markovian structure of the  $m$ -dimensional simple epidemic.

## 3. Main Results

Let  $(X_1(t), \dots, X_m(t))$  be an  $m$ -dimensional simple epidemic in a population consisting initially of  $b_i$  infectives from cause  $i$ ,  $i = 1, 2, \dots, m$ , and infinitely many susceptibles. This epidemic can be described by a sequence of bivariate random vectors,  $(T_1, \xi_1), (T_2, \xi_2), \dots, 2, \dots, T_1, T_2, \dots$ , are the interinfection times, and  $\xi_1, \xi_2, \dots$ , are random variables with values in  $\{1, \dots, m\}$ , which designate the respective infection causes. Let  $S_k$  be equal to  $\sum_{j=1}^k T_j$ ,  $k = 1, 2, \dots$ . The finite dimensional joint distributions of  $(X_1(t), X_m(t))$  are determined by  $(T_k, \xi_k)$ ,  $k = 1, 2, \dots$ , through the following set equalities:

$$(X_i(t) = b_i + k) = \bigcup_{r=k}^{\infty} (S_r \leq t < S_{r+1}, \sum_{q=1}^r I(\xi_q = i) = k) \quad (3.1)$$

for every positive real number  $t$ , positive integer  $k$ , and  $i = 1, \dots, m$ . We assume throughout that the sequence  $(T_1, \xi_1), (T_2, \xi_2), \dots$ , is determined by (3.2) and (3.3).

$$P\{\xi_k = \ell_k, k = 1, \dots, n\} =$$

$$\left[ \prod_{i=1}^m \prod_{j=0}^{r_i-1} (b_i + j) \right] \left[ \prod_{j=1}^n \left( \sum_{i=1}^m \alpha_i (b_i + \sum_{q=1}^{j-1} I(\ell_q = 1)) \right)^{-1} \right] \quad (3.2)$$

where  $r_i = \sum_{q=1}^n I(\ell_q = i)$ ,  $i = 1, \dots, m$ , and  $n = 1, 2, \dots$ .

$$P\{T_k | \xi_1, \dots, \xi_{k-1} > t_k, k = 1, \dots, n\} = e^{-\sum_{j=1}^n \mu_j t_j} \quad (3.3)$$

where  $t_1, \dots, t_n$  are positive real numbers,  $\mu_j = \sum_{i=1}^m \alpha_i (b_i + \sum_{q=1}^{j-1} I(\xi_q = 1))$ ,  $j = 1, \dots, n$ , and  $n = 1, 2, \dots$ .



We will show that for every two positive real numbers  $t$  and  $\beta$   $(X_{N,1}(t), \dots, X_{N,m}(t))$  converges in law to  $(X_1(t), \dots, X_m(t))$  as  $N \rightarrow \infty$ , and that  $\lim_{N \rightarrow \infty} E(X_{N,i}(t))^\beta = E(X_i(t))^\beta$  for  $i = 1, \dots, m$ . First we introduce two lemmas.

**Lemma 3.1.** Let  $U_1, U_2, \dots$ , be an i.i.d. sequence of exponential random variables with mean 1. Further let  $(\tau_{b+r-1,1}, \dots, \tau_{b+r-1,b+r-1})$  be the order statistic of a sample of size  $b+r-1$  from  $U_1$ . Then  $\sum_{j=1}^r (b+j-1)^{-1} U_j$  and  $\tau_{b+r-1,r}$  are identically distributed.

**Proof.** It suffices to note that the spacings;  $\tau_{r+b-1,r-j+1} - \tau_{r+j-1,r-j}$ ,  $j = 1, \dots, r$ , ( $\tau_{r+b-1,0} = 0$ ) are independent exponential random variables with means equal respectively to  $(b+j-1)^{-1}$ ,  $j = 1, \dots, r$ .

**Lemma 3.2.** Let  $t$  and  $\beta$  be positive real numbers; let  $c = \sup_N \sum_{i=1}^m b_{N,i}$  and  $\alpha = \max_{1 \leq i \leq m} \alpha_i$ . If (1.2) holds then; (i)  $P(S_{N,r} \leq t) \leq (1 - e^{-\alpha c t})^r$  for  $r = 1, \dots, N$ , and (ii)  $\sup_N E(X_{N,i}(t))^\beta < \infty$ , for  $i = 1, \dots, m$ .

**Proof.** We note that (ii) is a corollary of (i). To prove (i) let  $U_1, U_2, \dots$ , be as in Lemma 3.1. Since  $P(S_{N,k} \leq t) \leq P\{\sum_{j=1}^r j^{-1} U_j \leq \alpha c t\}$ , (i) follows from Lemma 3.1.

**Theorem 3.3.** Let  $\beta$  and  $t$  be positive real numbers. If (1.2) holds then;

(i)  $(X_{N,1}(t), \dots, X_{N,m}(t))$  converges in law as  $N \rightarrow \infty$  to  $(X_1(t), \dots, X_m(t))$ , and (ii)  $\lim_{N \rightarrow \infty} E(X_{N,i}(t))^\beta = E(X_i(t))^\beta$  for  $i = 1, \dots, m$ .

**Proof.** We note that (ii) is a corollary of (i) and Lemma 3.2 (ii).

To prove (i) let  $k_1, \dots, k_m$  be  $m$  nonnegative integers,  $A$  be the set  $\{(r_1, \dots, r_m) | r_i \geq k_i, i = 1, \dots, m\}$  and let  $R = \sum_{i=1}^m r_i$ . Since for  $N$

sufficiently large,  $P\{X_{N,i}(t) \geq k_i + b_i, i = 1, \dots, m\} = \sum_{A} P\{S_{N,R} \leq t < S_{N,R+1}, \sum_{q=1}^R I(\xi_{N,q} = i) = r_i, i = 1, \dots, m | R \leq N\}$ , (i) follows from Lemma (3.2) (i), the dominated convergence theorem, and some simple calculations.

For reference purposes the following two well known results are presented.

**Lemma 3.4.** Let  $U_1, U_2, \dots$ , be as in Lemma 3.1, and let  $\mu_1, \mu_2, \dots$ , be a sequence of positive real numbers. Further we denote by  $f_k$  the density function of  $\sum_{j=1}^k \mu_j^{-1} U_j$ ,  $k = 1, 2, \dots$ . Then, (i)  $\mu_{r+1}^{-1} f_{r+1}(t) = P\{\sum_{j=1}^r \mu_j^{-1} U_j \leq t < \sum_{j=1}^{r+1} \mu_j^{-1} U_j\}$ , and (ii)  $f_r(t) = (\prod_{j=1}^r \mu_j) \sum_{j=1}^r e^{-\mu_j t} \prod_{i=1, i \neq j}^r (\mu_i - \mu_j)^{-1}$ , provided  $\mu_1, \dots, \mu_r$  are distinct.

We are ready to state and prove the main result of the paper.

**Theorem 3.5.** Let  $t$  and  $\beta$  be positive real numbers, and  $k_1, \dots, k_m$  be nonnegative integers. Further let  $A$  be the set  $\{(r_1, \dots, r_m) | r_i \geq k_i, i = 1, \dots, m\}$ . If conditions (1.1) and (1.2) are satisfied, then the following hold.

- (i)  $\lim_{N \rightarrow \infty} P\{X_{N,i}(t) \geq k_i + b_i, i = 1, \dots, m\} = \sum_A \prod_{i=1}^m \binom{b_i + r_i - 1}{b_i - 1} e^{-\alpha_i b_i t} (1 - e^{-\alpha_i t})^{k_i}$ .
- (ii)  $\lim_{N \rightarrow \infty} E(X_{N,i}(t))^\beta$  is the  $\beta$ th moment of a negative binomial random variable with parameters  $b_i$  and  $e^{-\alpha_i t}$ ,  $i = 1, \dots, m$ .

Proof. By Theorem 3.3 it suffices to show that  $X_1(t), \dots, X_m(t)$  are independent, and have the respective negative binomial marginal distributions.

Let  $k_1, \dots, k_m$  be nonnegative integers, with sum equal to  $k$ . Further, let  $B$  denote the set

$$[(\ell_1, \dots, \ell_k) | \ell_1, \dots, \ell_k \in \{1, \dots, m\}, \sum_{q=1}^k I(\ell_q = i) = k_i, i = 1, \dots, m].$$

We note that  $P\{X_i(t) = k_i + b_i, i = 1, \dots, m\}$  equals

$$\sum_B P\{S_k \leq t < S_{k+1} | \xi_q = \ell_q, q = 1, \dots, k\} P\{\xi_q = \ell_q, q = 1, \dots, k\}.$$

From Lemma 3.4 and equation (3.2) it follows that  $P\{X_i(t) = k_i + b_i, i = 1, \dots, m\}$  equals

$$\prod_{i=1}^m \binom{b_i + k_i - 1}{b_i - 1} e^{-\alpha_i b_i t} \frac{\alpha_i^{k_i} k_i!}{\alpha_i^{k_i} k_i!} \sum_B \sum_{j=1}^{k+1} e^{-\theta_j t} \frac{j^{k+1}}{\prod_{\substack{i=1 \\ i \neq j}}^{k+1} (\theta_i - \theta_j)^{-1}}, \text{ where}$$

$$\theta_j = \sum_{q=1}^{j-1} \sum_{i=1}^m \alpha_i I(\ell_q = i), j = 1, \dots, k+1.$$

By reusing Lemma 3.4 we obtain that  $P\{X_i(t) = k_i + b_i, i = 1, \dots, k\}$  equals

$$\prod_{i=1}^m \binom{b_i + k_i - 1}{b_i - 1} e^{-\alpha_i b_i t} \frac{\alpha_i^{k_i} k_i!}{\alpha_i^{k_i} k_i!} \sum_B P\left\{\sum_{j=1}^{k+1} \theta_j^{-1} U_j \leq t\right\} \prod_{j=2}^{k+1} \theta_j^{-1}.$$

To complete the proof it suffices to show that

$$\prod_{i=1}^m \alpha_i^{k_i} k_i! \sum_B P\left\{\sum_{j=2}^{k+1} \theta_j^{-1} U_j \leq t\right\} \prod_{j=2}^{k+1} \theta_j^{-1} = \prod_{i=1}^m (1 - e^{-\alpha_i t})^{k_i}. \quad (3.4)$$

Let  $F$  denote the left side of equation (3.4). Further let

$$B_i = [(\ell_1, \ell_2, \dots, \ell_q) | \ell_2, \dots, \ell_k \in \{1, \dots, m\}, \sum_{q=2}^k I(\ell_q = r) = k_r, r = 1, \dots, m, r \neq i, \sum_{q=2}^k I(\ell_q = i) = k_i - 1].$$



We observe that  $F(t)$  equals  $\prod_{i=1}^m \alpha_i^{k_i} k_i! \sum_{i=1}^m \sum_{B_i} P\{\sum_{j=2}^{k+1} \theta_j^{-1} U_j \leq t\} \prod_{j=2}^{k+1} \theta_j^{-1}$ .

By a simple calculation and Lemma 3.4, it follows that

$$\frac{dF}{dt} = \prod_{i=1}^m \alpha_i^{k_i} k_i! \sum_{i=1}^m e^{-\alpha_i t} \sum_{B_i} P\{\sum_{j=3}^{k+1} \theta_{j,i}^{-1} U_j \leq t\} \prod_{j=3}^{k+1} \theta_{j,i}^{-1}, \text{ where } \theta_{j,i} = \sum_{\substack{r=1 \\ r \neq i}}^m \alpha_r \sum_{q=2}^{j-1} I(l_q = r), \quad i = 1, \dots, m, \quad j = 3, \dots, k+1.$$

Consequently Equation

(3.4) is established by an induction argument on  $k$  applied to each of the sums  $\sum_{B_i} P\{\sum_{j=3}^{k+2} \theta_{j,i}^{-1} U_j \leq t\} \prod_{j=3}^{k+1} \theta_{j,i}^{-1}, \quad i = 1, \dots, m.$

Corollary 3.6. Let  $X_N(t)$  the total number of infectives at time  $t$ . If

(1.1) and (1.2) are satisfied, then the following hold.

- (i)  $X_N(t)$  converges in law as  $N \rightarrow \infty$  to a sum of  $m$  independent negative binomial random variables with respective parameters  $b_i, e^{-\alpha_i t}, i = 1, \dots, m$ , and
- (ii)  $\lim_{N \rightarrow \infty} E(X_N(t))^\beta$  equals to the  $\beta$ th moment of the limiting random variable given in (i).

In the particular case discussed in Billard, Lacayo and Langberg (1978),

$X_N(t)$  converges in law to a negative binomial random variable with parameters

$\sum_{i=1}^m b_i$  and  $e^{-\alpha t}$ . This fact played a major role in the proof of (1.3) for

the symmetric case as presented in the cited reference. Finally we note

that it is of interest to investigate the asymptotic behavior of  $m$ -dimensional simple epidemics, that do not have the Markovian structure.

## REFERENCE

- (1) Billard, L., Lacayo, H. and Langberg, Naftali A. (1978). The Symmetric  $m$ -dimensional Simple Epidemic. FSU Statistics Report M461.

## UNCLASSIFIED

## SECURITY CLASSIFICATION OF THIS PAGE

## REPORT DOCUMENTATION PAGE

1. REPORT NUMBER FSU No. M464 AFOSR No. 84 AFOSRN No. 9	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE On the Negative Binomial Convergence in a Class of M-Dimensional Simple Epidemics	5. TYPE OF REPORT & PERIOD COVERED Technical Report	6. PERFORMING ORG. REPORT NUMBER FSU Statistics Report M464
7. AUTHOR(s) H. Lacayo Naftali A. Langberg	8. CONTRACT OR GRANT NUMBER(s) AFOSR 74-2581D AFOSR 76-3109	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
9. PERFORMING ORGANIZATION NAME & ADDRESS The Florida State University Department of Statistics Tallahassee, Florida 32306	11. CONTROLLING OFFICE NAME & ADDRESS The U.S. Air Force Air Force Office of Scientific Research Bolling Air Force Base, D.C. 20332	12. REPORT DATE July, 1978
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	13. NUMBER OF PAGES 10	15. SECURITY CLASS (of this report) Unclassified
16. DISTRIBUTION STATEMENT (of this report) Approved for public release; distribution unlimited.		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS Exponential distributions, convergence in law, stochastic processes, m-dimensional simple epidemics, negative binomial distributions		
20. ABSTRACT		

We consider a population which is exposed to  $m$  infections, and consist initially of  $N$  susceptibles. At each point in time at most one susceptible becomes infective, and only from one cause. This  $m$ -dimensional simple epidemic is a stochastic process,  $(X_{N,1}(t), \dots, X_{N,m}(t))$ , with components counting the number of infectives from the respective causes at time  $t$ . We show that if the transition rates of cause 1 through  $m$  at time  $t$  are given by

$$\alpha_i X_{N,i}(t) \left[ 1 - \frac{1}{N} \sum_{i=1}^m (X_{N,i}(t) - X_{N,i}(0)) \right], i = 1, \dots, m, \alpha_1, \dots, \alpha_m > 0 \text{ and if } \lim_{N \rightarrow \infty} X_{N,i}(0) = b_i \in \{1, 2, \dots\},$$

then  $(X_{N,1}(t), \dots, X_{N,m}(t))$  converges as  $N \rightarrow \infty$  to a random vector with independent negative binomial components.